

AD-A171 577


NMR (NUCLEAR MAGNETIC RESONANCE) INVESTIGATIONS OF
LIQUID PROPELLANT SYST (U) ARMY ARMAMENT RESEARCH AND
DEVELOPMENT CENTER DOVER NJ ARMANE S BULUSU ET AL
SEP 86 ARAED-TR-86028


1/1

UNCLASSIFIED

F/G 19/1

NL





PHOTOCOPY RESOLUTION TEST
NATIONAL BUREAU OF STANDARDS

AD-A171 577

12
AD

AD-E401 556

TECHNICAL REPORT ARAED-TR-86028

**NMR INVESTIGATIONS OF LIQUID PROPELLANT SYSTEMS
CONSISTING OF ALKYL- AND HYDROXYL-AMMONIUM NITRATES**

**PART 1. CHARACTERIZATION OF CARBON-13, PROTON, AND NITROGEN-15
NMR SPECTRA AND ANALYTICAL APPLICATIONS**

S. BULUSU
R. L. DUDLEY

DTIC
ELECTE
SEP 4 1986
S A D

SEPTEMBER 1986



ARMAMENT RDE CENTER

U. S. ARMY ARMAMENT RESEARCH, DEVELOPMENT AND ENGINEERING CENTER

ARMAMENT ENGINEERING DIRECTORATE

DOVER, NEW JERSEY

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

DTIC FILE COPY

86

9

04

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

Destroy this report when no longer needed by any method that will prevent disclosure of contents or reconstruction of the document. Do not return to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report ARAED-TR-86028	2. GOVT ACCESSION NO ADA171577	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NMR INVESTIGATIONS OF LIQUID PROPELLANT SYSTEMS CONSISTING OF ALKYL- AND HYDROXYL-AMMONIUM NITRATES PART I. CHARACTERIZATION OF CARBON-13, PROTON, AND NITROGEN-15 NMR SPECTRA AND ANALYTICAL APPLICATIONS		5. TYPE OF REPORT & PERIOD COVERED Final
6. PERFORMING ORG. REPORT NUMBER		7. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS ARDEC, AED Energetics & Warheads Div (SMCAR-AEE) Dover, NJ 07801-5001		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 1L161102AH60
11. CONTROLLING OFFICE NAME AND ADDRESS ARDEC, IMD STINFO Div (SMCAR-MSI) Dover, NJ 07801-5001		12. REPORT DATE September 1986
14. MONITORING AGENCY NAME & ADDRESS (If different from Controlling Office)		13. NUMBER OF PAGES 25
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This project was initiated in 1985 by the Large Caliber Weapon Systems Laboratory, Energetic Materials Division, ARDC.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Liquid propellants Hydroxyl ammonium nitrate (HAN) NMR spectroscopy Oxidizers Aliphatic ammonium nitrates (AAN's) Monopropellant fluids		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The proton, carbon-13, and nitrogen-15 (natural abundance) nuclear magnetic resonance (NMR) spectra of four aliphatic ammonium nitrates (AAN's) which could find application as the fuel components in liquid propellant systems and the nitrogen-15 NMR spectrum of hydroxyl ammonium nitrate (HAN) which serves as the oxidizer have been characterized in terms of their chemical shifts. (continued)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT (Continued)

The utility of NMR spectroscopy for analytical applications has been evaluated. It was found that while the AAN's can be analyzed by using either proton or carbon-13 NMR, NMR spectroscopy cannot be applied readily to the analysis of HAN. Nitrogen-15 NMR at natural abundance may be applied to analyze HAN, but the technique would be too time-consuming.

UNCLASSIFIED

• • SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE OF CONTENTS

	Page
Introduction	1
Experimental	2
¹ H Spectra	2
¹³ C Spectra	2
¹⁵ N Spectra	
Results and Discussion	3
Proton (¹ H) NMR Spectra	3
¹³ C-NMR Spectra	3
¹⁵ N-NMR Spectra	4
Conclusions	4
References	17
Distribution List	19



A-1

TABLE

	Page
1 Summary of the proton chemical shifts δ of the AAN's: IPAN, TMAN, TEAN, and NBAN	5

FIGURES

1 ^1H -NMR spectrum of IPAN in D_2O solution	7
2 ^1H -NMR spectrum of TMAN in D_2O solution	8
3 ^1H -NMR spectrum of TEAN in D_2O solution	9
4 ^1H -NMR spectrum of NBAN in D_2O solution	10
5 Carbon-13 NMR spectra of aqueous solutions of TMAN (1.8 moles/l), TEAN (1.07 moles/l), and IPAN (1.8 moles/l)	11
6 Carbon-13 NMR spectrum of NBAN in H_2O	12
7 Plots of concentration versus peak heights for IPAN	13
8 Plots of concentration versus peak heights for TEAN and TMAN	14
9 ^1H -coupled C-13 spectrum of IPAN. $J(^{13}\text{C}-^1\text{H})$: CH_3 , 125.8 Hz; CH , 143.3 Hz	15
10 Proton decoupled nitrogen 15 NMR spectra of:	16
A) Nitrogen-15 labelled AN in H_2O with a D_2O lock. Delay time (D_1) 60 secs; number of transients, (NT) 16; acquisition time (AT), 0.5 sec.	
B) Hydroxyl ammonium nitrate (80%) in H_2O with a D_2O lock. D_1 , 15 secs; NT, 500; AT, 0.5 sec.	
C) Isopropyl AN in DMSO with a D_2O lock, D_1 , 10 secs; NT, 4456; AT, 0.5 secs.	
D) n-butyl AN in DMSO with a D_2O lock, D_1 , 10 secs; NT, 3900; AT, 0.5 secs.	
E) Triethanol AN in DMSO with a D_2O lock. D_1 15 secs; NT, 5000; AT, 0.5 secs.	
F) Trimethyl AN in DMSO with D_2O lock. D_1 , 15 secs; NT, 15, 224; AT, 0.5 secs.	

INTRODUCTION

Liquid propellants which consist of hydroxyl ammonium nitrate (HAN), an aliphatic ammonium nitrate (AAN) and just sufficient water to keep them fluid over a wide temperature range, have been under investigation for some time to substitute solid propellants in guns (refs 1-4). These "monopropellant" fluids which contain both the fuel and the oxidizer for self-sustained combustion, offer high volumetric impetus and some important advantages of vehicle space utilization and improved mobility. However, impurities associated with the amine nitrates, especially metal ions (e.g., Cu^{2+} and Fe^{3+}) seem to adversely affect their ballistic performance and stability in general, calling for the use of sequestering agents to eliminate the metal ions. Methods of achieving and monitoring the purity and concentration of the liquid propellants during the production and storage are therefore in need of development.

Typically, the liquid propellants contain 80 weight percent or more of salt in aqueous solution and have, therefore, to be considered as extremely concentrated electrolytic solutions which may be compared to molten salts (ref 1) in many properties. The microstructure of these "liquids," that is, the ionic interactions and the coordination chemistry among the various molecular and ionic species present in them are poorly understood, but are of obvious importance to their proper application and prediction of performance. The present nuclear magnetic resonance (NMR) investigation was initiated for the dual purpose of exploring the potential use of NMR spectroscopy for analytical applications and studying the microstructure and ionic interactions mentioned above in the highly concentrated, ionic, liquid propellant systems. This Part I of the report deals with the characterization of the NMR spectra for analytical applications and the Part II addresses the ionic microstructure of the liquid propellants selected for the study.

Three AAN's were previously selected as prime candidates (ref 1) for use as fuels in the HAN-based liquid propellants based on their thermal behavior and detonation susceptibility. These were isopropyl ammonium nitrate (IPAN), trimethyl ammonium nitrate (TMAN), and triethanol-amine nitrate (TEAN). While these three AAN's were the primary focus of this investigation, n-butylammonium nitrate (NBAN) is also included since much information on the carbon-13 spin-relaxation times (T_1 , s) of several salts of n-butylamine relevant to Part II of this investigation, is available in the literature.

Previous analytical approaches (ref 5) to liquid propellants consisted of UV spectrophotometric and conductometric methods for the determination of NO_3^- and hydroxylamine and atomic absorption spectrometry for the parts per million level of transition metal ion impurities. Only a cursory attempt (ref 4) seemed to have been made to apply proton (^1H)-NMR to the analysis of the AAN's. In this report (Part I) the hydrogen-1 and the natural abundance carbon-13 (^{13}C) and nitrogen-15 (^{15}N) NMR spectra are presented with a discussion of their analytical usefulness.

EXPERIMENTAL

All the NMR spectra included in this report were obtained on a Varian Associates XL-200 pulse fourier transform spectrometer equipped with a proton probe and a separate broad-band probe for ^{13}C and ^{15}N . The spectrometer was operated with an internal D_2O lock for proton spectra and an external D_2O lock for the ^{13}C and ^{15}N spectra, respectively.

^1H Spectra

The proton spectra were obtained with approximately 50 mg of the AAN dissolved in 0.5 mL D_2O using 5 mm sample tubes. Typical acquisition parameters were: pulse width (pw) = 6-8 μsec , acquisition time (AT) = 2 sec, D_1 (pulse interval) = 10 sec, and NT (number of transients) \approx 10. The spectra were referenced using 4,4-dimethyl-4-silapentane sodium sulfonate (DDS) ($\delta^1\text{H}$ = 0.015 ppm) as an internal standard.

^{13}C Spectra

For obtaining the ^{13}C spectra 0.5 g each of the AAN's was dissolved in 2.2 mL H_2O contained in a 10 mm NMR sample tube. A 5 mm coaxial tube filled with D_2O was inserted into the 10 mm sample tubes to serve as external D_2O lock. The following acquisition parameters were employed based on an approximate determination of the Ernst angle: pw = 5 μsec , AT = 2 sec, D_1 = 0 and NT \approx 250. Adequate signal to noise ratio can in fact be obtained with a considerably smaller number of transients. The chemical shift scale was calibrated using $\delta^{13}\text{C}$ = 132.1 ppm (relative to TMS) for the aromatic carbons of hexamethyl benzene as an external standard.

^{15}N Spectra

In this preliminary work attempts were made to obtain ^{15}N -NMR spectra at natural abundance. However, good signal-to-noise ratios were difficult to obtain particularly for the NO_3 -nitrogen. Ammonium- ^{15}N signals could be obtained with reasonable signal-to-noise ratios using long (>30 sec) pulse intervals and DMSO-d_6 , as solvent.

RESULTS AND DISCUSSION

Proton (^1H) NMR Spectra

The proton NMR spectra of the AAN's, IPAN, TMAN, TEAN, and NBAN are shown in figures 1 to 4. In common to all the four AAN's, the H_2O peak derived from contamination in D_2O and exchange with the labile protons can be seen at ~ 4.8 ppm. The labile protons attached to the cationic nitrogen would be highly deshielded and may be observed only in a nonaqueous solvent. This fact was independently confirmed but not shown in the figures. The protons attached to the carbons in each of the salts can be readily assigned as indicated in the spectra using the spin-coupling information. All the proton chemical shift (δ , ppm) information is summarized in table 1 for convenience.

Since only one AAN will be present along with HAN in any given liquid propellant system, the use of the above proton peak areas is not expected to present any problem in quantitative analysis. However, HAN cannot be analyzed by this method since all the protons in this structure would be exchangeable and labile. The concentration of the protons attached to the carbons can be measured without interference from HAN.

^{13}C -NMR Spectra

The proton noise decoupled ^{13}C spectra of the four AAN's, IPAN, TMAN, TEAN, and NBAN, are shown in figures 5 and 6, respectively. In these spectra each chemically distinguishable carbon gives rise to a separate line readily assignable on the basis of the well known shielding and inductive effects. The assignments are indicated in the figures.

It may be pointed out that the peak intensities in carbon-13 NMR spectra do not necessarily bear an exact relationship to the number of carbons of each type in the structure. This is due to the interplay of two effects. Firstly, the large and widely varying relaxation (T_1) times of different carbon-13 nuclei compared to the protons, requiring different pulse widths and pulse intervals to avoid peak saturation. Secondly, the inherent low sensitivity of carbon-13 nucleus (1.7×10^{-4} relative to ^1H) commonly requires proton noise decoupling to enhance the sensitivity. The enhancement in sensitivity, up to a maximum of 3 times, is brought about by nuclear overhauser effect (NOE). These two effects make it often difficult to relate the peak intensities in a spectrum to the number of each type of carbon atoms in the structure under investigation. However, the peak intensity of any given carbon in the structure chosen for quantitative analysis can be optimized to maintain a linear relationship to the concentration of that substance in a fixed volume of solution. This is illustrated in figure 7 representing a plot of concentration versus peak height for the primary and the tertiary carbons in IPAN and in figure 8 for the carbons of TEAN and TMAN. For peaks which are considerably broader peak areas are preferable for quantitative analysis. This relationship was again found to be unaffected by the presence of HAN.

The proton noise decoupling which gives rise to the NOE also results in a simplification in the carbon-13 spectra which can be difficult to interpret otherwise. The strong proton couplings, such as those illustrated in figure 9 for IPAN, give rise to a multitude of overlapping lines in more complex structures.

¹⁵N-NMR Spectra

Nitrogen-15 NMR spectra at natural abundance (~0.35%) are often difficult to obtain especially when the spin-lattice relaxation times are long. The AAN's studied thus far gave poor signal-to-noise ratios even with pulse intervals of the order of 30 sec. Detectable signals from the ammonium nitrogen can be observed more readily owing to their shorter relaxation times resulting from the attached protons. The natural abundance ¹⁵N spectra of ¹⁵NH₄⁺, ¹⁵NO₃⁻, IPAN, TMAN, TEAN, and NBAN are collected together in figure 10. The acquisition parameters are described in the respective figure captions. The ¹⁵N- chemical shifts of the ammonium nitrogens in the AAN's taken from these spectra are summarized below. In these spectra, ¹⁵NH₄⁺, ¹⁵NO₃⁻, IPAN, TEAN, and NBAN show ¹⁵NO₃⁻ absorptions also in addition to the ammonium nitrogen lines. The higher noise levels in some of the spectra can be readily improved by accumulating more transients. However, because of the long delay times required and the NOE effects on nitrogen absorptions it would be impractical to use these spectra for purely analytical applications.

AAN in DMSO-d ₆ Solution	$\delta(^{15}\text{N}^+)$, ppm from NO ₃ -nitrogen
IPAN	-328.8
TEAN	-330.3
TMAN	-345.3
NBAN	-343.3
HAN	-292.5


Attempts will be made in the future to synthesize the AAN's enriched with ¹⁵N in both ammonium and nitrate nitrogens in order to characterize the N-15 spectra better and investigate the spin-lattice relaxation times (see Part II of this report).

CONCLUSIONS

The ¹H and ¹³C NMR spectra of the four AAN's studied may be readily adapted for the analysis of these materials in the liquid propellant systems. The natural abundance ¹⁵N spectra of the AAN's and the HAN, however, take typically 15 to 20 hours each, to yield reasonable signal-to-noise ratios and hence are judged impractical for analytical applications. It is concluded, therefore, that the HAN cannot readily be analyzed using NMR spectroscopy while the AAN's can be analyzed by either ¹Hp or ¹³C-NMR.


Table 1. Summary of the proton chemical shifts δ of the AAN's:
IPAN, TMAN, TEAN, and NBAN

	$\delta^1\text{H}$, ppm (TMS = 0)	Hz
TEAN in D ₂ O		
Downfield triplet	3.99	789
	3.97	793
	3.94	788
Upfield triplet	3.52	704
	3.49	699
	3.47	694
H ₂ O in D ₂ O	4.79	959
DSS	0.02	3
IPAN in D ₂ O		
H ₂ O in D ₂ O	4.81	963
Center of septet	3.50	699
CH ₃ doublet	1.32	253
	1.28	257
TMAN in D ₂ O		
H ₂ O in D ₂ O	4.88	976
CH ₃	2.88	576
NBAN in D ₂ O		
H ₂ O in D ₂ O	4.77	955
¹ C-triplet	2.98	596
² C-quintet	1.62	324
³ C-center doublet	1.38, 1.34	276, 268
⁴ C-triplet	0.90	181




^1H -NMR Spectrum
 $(\text{CH}_3)_2\text{CH}\text{-NH}^+\text{NO}_3^-$

Figure 1. ^1H -NMR spectrum of IPAN in D_2O solution



^1H -NMR Spectrum
 $(\text{HOCH}_2\text{CH}_2)_3\text{NH}^+\text{NO}_3^-$

Figure 2. ^1H -NMR spectrum of TMAN in D_2O solution



^1H -NMR Spectrum
 $(\text{CH}_3)_3\text{NH}^+\text{NO}_3^-$

Figure 3. ^1H -NMR spectrum of TEAN in D_2O solution






Figure 4. $^1\text{H-NMR}$ spectrum of NBAN in D_2O solution




^{13}C -NMR Spectra

Figure 5. Carbon-13 NMR spectra of aqueous solutions of TMAN (1.8 moles/l), TEAN (1.07 moles/l), and IPAN (1.8 moles/l)



^{13}C -NMR Spectrum
 $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_3^+\text{NO}_3^-$

Figure 6. Carbon-13 NMR spectrum of NBAN in H_2O



Concentration vs Peak Heights
(CH₃)₂CH-NH₃⁺NO₃⁻ (IPAN)

Figure 7. Plots of concentration versus peak heights for IPAN





Figure 8. Plots of concentration versus peak heights for TEAN and TMAN



^1H -Coupled ^{13}C -Spectrum of IPAN
 $(\text{CH}_3)_2\text{CHNH}_3^+\text{NO}_3^-$

Figure 9. ^1H -coupled C-13 spectrum of IPAN. $J(^{13}\text{C}-^1\text{H})$: CH_3 , 125.8 Hz;
 CH , 143.3 Hz




Figure 10. Proton decoupled nitrogen 15 NMR spectra of: A) Nitrogen-15 labeled AN in H_2O with a D_2O lock. Delay time (D_1) 60 secs; number of transients, (NT) 16; acquisition time (AT), 0.5 sec. B) Hydroxylammonium nitrate (80%) in H_2O with a D_2O lock. D_1 , 15 secs; NT, 500; AT, 0.5 sec. C) Isopropyl AN in DMSO with a D_2O lock, D_1 , 10 secs; NT, 4456; AT, 0.5 secs. D) n-butyl AN in DMSO with a D_2O lock, D_1 , 10 secs; NT, 3900; AT, 0.5 secs. E) Triethanol AN in DMSO with a D_2O lock. D_1 , 15 secs; NT, 5000; AT, 0.5 secs. F) Trimethyl AN in DMSO with D_2O lock. D_1 , 15 secs; NT, 15, 224; AT, 0.5 secs.

REFERENCES

1. N. Klein, CPIA Publication 292, Vol II, 14th JANNAF Combustion Meeting, 1977.
2. N. Klein, "Preparation and Characterization of Several Liquid Propellants," Technical Report ARBRL-TR-02471, Ballistic Research Laboratory, Aberdeen, MD, 1983.
3. N. Klein, "Isomer Effects on Liquid Propellant Performance," 19th JANNAF Combustion Meeting, CPIA Publication No. 366, Vol I, p. 513, 1982.
4. N. Klein, K.E. Travis, and R.A. Sasse, "Methods for Product Analysis in Production of Hydroxyl-Ammonium Nitrate Propellants," International Symposium, Analytical Methods for Propellants and Explosives, 22-24 June 1977, p. 325.
5. N. Klein, K.E. Travis, and R.A. Sasse, "Methods for Product Analysis in Production of Hydroxylammonium Nitrate Propellants," International Annual Conference on Analytical Methods for Propellants and Explosives, Karlsruhe, W. Germany, 1977.
6. "Practical NMR Spectroscopy," by M.L. Martin, J.J. Depuech, and G.L. Martin, Heydon & Sons, Ltd., London, 1980.

DISTRIBUTION LIST

Commander

Armament Research, Development and
Engineering Center

ATTN: SMCAR-AE, R. Bushey

J.P. Picard

SMCAR-AEE, J.A. Lannon

Word Processing Office (3)

SMCAR-AEE-B, D. Downs

SMCAR-AEE-BR, L. Harris

A. Beardell

B. Brodman

Y. Carignan

A. Bracuti

W. Seals

SMCAR-AEE-WW, C. Capellos

F. Owens

P. Lu

S. Nicolaides

SMCAR-AER-T, P. Marinkas

SMCAR-MSI (5)

SMCAR-TD

SMCAR-TDC, H. Grundler

Dover, NJ 07801-5001

Commander

U.S. Army Armament, Munitions and Chemical Command

ATTN: AMSMC-GCL(D)

Dover, NJ 07801-5001

Administrator

Defense Technical Information Center

ATTN: Acquisitions Division

Cameron Station

Alexandria, VA 22304-6145

Director

U.S. Army Materiel Systems Analysis Activity

ATTN: AMXSY-MP

Aberdeen Proving Ground, MD 21005-5066

Commander

Chemical Research, Development
and Engineering Center

U.S. Army Armament, Munitions and Chemical Command

ATTN: SMCCR-SPS-IL

Aberdeen Proving Ground, MD 21010-5423

Commander
Chemical Research, Development
and Engineering Center
U.S. Army Armament, Munitions and Chemical Command
ATTN: SMCCR-RSP-A
Aberdeen Proving Ground, MD 21010-5423

Director
Ballistic Research Laboratory
ATTN: AMXBR-OD-ST
Aberdeen Proving Ground, MD 21005-5066

Chief
Benet Weapons Laboratory, CCAC
Armament Research, Development and Development Center
U.S. Army Armament, Munitions and Chemical Command
ATTN: SMCAR-CCB-TR
Watervliet, NY 12189-5000

Commander
U.S. Army Armament, Munitions and Chemical Command
ATTN: SMCAR-ESP-L
Rock Island, IL 61299-6000

Director
U.S. Army TRADOC Systems Analysis Activity
ATTN: ATAA-SL
White Sands Missile Range, NM 88002

HDQA
ATTN: DAMA-ART-M
DAMA-CSM
DAMA-ZA
Washington, DC 20310

Commander
U.S. Army Materiel Command
ATTN: AMCPM-GCM-WF
5001 Eisenhower Avenue
Alexandria, VA 22304

Commander
U.S. Army Materiel Command
ATTN: AMCDRA-ST
5001 Eisenhower Avenue
Alexandria, VA 22333-0001

Project Manager
Tank Main Armament Systems
ATTN: AMCPM-TMA, K. Russell - 120
AMCPM-TMA-105
AMCPM-TMA-120
Dover, NJ 07801-5001

Director
Ballistic Research Laboratory
ATTN: SMCAR-BR-IB-B, N. Klein
E. Friedman
W.R. Morrison
A. Miziolek
Aberdeen Proving Ground, MD 21005-5066

Commander
LABCOM
ATTN: AMSLC-TP-PL, R. Vitali
I. Bartky
2800 Powder Mill Road
Adelphi, MD 20783-1145

Director of Army Research and Technology
ATTN: DAMA-AR, L.M. Cameron
Pentagon - Room 3E474
Washington, DC 20310-0630

END

DTIC

10-86